

Stratification of Moving Multicomponent Bose-Einstein Condensates

V.I. Yukalov¹ and E.P. Yukalova²

¹*Bogolubov Laboratory of Theoretical Physics
Joint Institute for Nuclear Research, Dubna 141980, Russia*

²*Department of Computational Physics
Laboratory of Information Technologies
Joint Institute for Nuclear Research, Dubna 141980, Russia*

Abstract

A mixture of the multicomponent Bose-Einstein condensate is considered, where each component moves with its own velocity. As a result of the relative motion, the mixture stratifies when the relative velocity reaches a critical value. Stability conditions for a binary moving mixture are derived and the critical velocity is found.

Ultracold atomic gases, forming Bose-Einstein condensates, are nowadays an active field of research (see reviews [1–5]). Dilute Bose gas can be trapped and cooled down to temperatures when practically all atoms are in Bose-Einstein condensate. One can create mixtures of several Bose-condensed species. The stability of such mixtures depends on the mutual atomic interactions. The condition of stability for binary mixtures was, first, found by Nepomnyashchy [6], considering the collective excitation spectrum in the Bogolubov and long-wave approximations. When this stability condition is not valid, the mixture is not stable and stratifies onto separate components. The analysis of Nepomnyashchy [6] was based on the Green functions approach for a superfluid mixture with broken gauge symmetry. As was shown later [7], the same stability condition holds true for a fluid mixture with conserved gauge symmetry.

Recently, it has become feasible to create moving wave packets of Bose-Einstein condensates, which could collide and move through each other [8–12]. Therefore, it is interesting to study the interplay between the possibility of different Bose components to mix and the influence on their stability of their relative motion through each other. Such a relative motion of two fluids could change the stability condition of mixing [6,7]. Bose-Einstein condensates, formed in modern traps, are made of dilute gases, for which the Gross-Pitaevskii equation provides a very good description [1–5]. Actually, this equation is an exact equation for the coherent system at zero temperature [13]. The aim of the present communication is to analyze a mixture of Bose-Einstein condensates, whose components move through each other, under conditions of low temperature, when the system is described by the Gross-Pitaevskii equation. This situation corresponds to current experiments with Bose-Einstein condensates of dilute gases.

An atomic gas, of N atoms in volume V , is dilute when its density $\rho \equiv N/V$ and the s -wave scattering length a are connected by the inequality $\rho a^3 \ll 1$. This can be generalized to a mixture of species enumerated by the index $i = 1, 2, \dots$ as the inequality $\sqrt{\rho_i \rho_j} a_{ij}^3 \ll 1$, in which $a_{ii} \equiv a_i$ and a_{ij} are the corresponding scattering lengths, and $\rho_i \equiv N_i/V$. The Gross-Pitaevskii equation for a multicomponent mixture is

$$i\hbar \frac{\partial}{\partial t} \eta_j(\mathbf{r}, t) = \hat{H}_j[\eta] \eta_j(\mathbf{r}, t) , \quad (1)$$

where $\eta_i(\mathbf{r}, t)$ is the coherent field of the i -component; in the nonlinear Hamiltonian

$$\hat{H}_i[\eta] = \frac{\hat{\mathbf{p}}^2}{2m_i} + U_i + \sum_j A_{ij} |\eta_j|^2 \quad (2)$$

the first term is the kinetic-energy operator, with $\hat{\mathbf{p}} \equiv -i\hbar \vec{\nabla}$, and m_i being atomic mass, U_i is an external (trapping) potential, and the third term describes contact atomic interactions with

$$A_{ij} \equiv 2\pi\hbar^2 a_{ij} \frac{m_i + m_j}{m_i m_j} .$$

The normalization condition for each component reads

$$||\eta_i||^2 \equiv \int |\eta_i(\mathbf{r}, t)|^2 d\mathbf{r} = N_i .$$

Elementary collective excitations of the system can be obtained by linearizing the Gross-Pitaevskii equation (1) around a given stationary solution. Following here this way, we should keep in mind that our aim here is to study a mixture of moving components. Assume that each component moves with a fixed velocity $\mathbf{s}_i$. Then taking account of this motion can be done by means of the Galilean transformation replacing η_j by $\eta_j \exp\{\frac{i}{\hbar} m_j \mathbf{s}_j \cdot \mathbf{r}\}$. To find collective excitations in the moving mixture, we write the coherent field η_j in the form

$$\eta_j(\mathbf{r}, t) = [\varphi_j(\mathbf{r}) + u_j(\mathbf{r})e^{-i\omega t} + v_j^*(\mathbf{r})e^{i\omega t}] \exp\left\{\frac{i}{\hbar} (m_j \mathbf{s}_j \cdot \mathbf{r} - E_j t)\right\}, \quad (3)$$

where $\varphi_j(\mathbf{r})$ satisfies the stationary equation

$$\left(\hat{H}_i[\varphi] + \hat{\mathbf{p}} \cdot \mathbf{s}_i + \frac{m_i s_i^2}{2}\right) \varphi_i(\mathbf{r}) = E_i \varphi_i(\mathbf{r}), \quad (4)$$

with the normalization $||\varphi_i||^2 = N_i$. Substituting Eq. (3) into the Gross-Pitaevskii equation (1) and linearizing it with respect to the functions $u_j(\mathbf{r})$ and $v_j(\mathbf{r})$, we obtain

$$\begin{aligned} \left(\hbar\omega - \hat{\mathbf{p}} \cdot \mathbf{s}_i - \hat{H}[\varphi] - \frac{m_i s_i^2}{2} + E_i\right) u_i - \varphi_i \sum_j A_{ij} (\varphi_j^* u_j + \varphi_j v_j) &= 0, \\ \left(\hbar\omega - \hat{\mathbf{p}} \cdot \mathbf{s}_i + \hat{H}[\varphi] + \frac{m_i s_i^2}{2} - E_i\right) v_i + \varphi_i^* \sum_j A_{ij} (\varphi_j^* u_j + \varphi_j v_j) &= 0. \end{aligned} \quad (5)$$

This is a generalization of the Bogolubov - de-Gennes equations to the case of a mixture of moving components.

Being interested in collective excitations above the ground state of trapped atoms, we keep in mind that the wave functions, defined by Eq. (4), are real-valued. Then we can resort to the local-density approximation, setting $\varphi_i(\mathbf{r}) \simeq \sqrt{\rho_i(\mathbf{r})}$ and assuming that u_i and v_i are proportional to $e^{i\mathbf{k} \cdot \mathbf{r}}$. Supposing that atomic interactions are sufficiently strong, so that mean potential energies are essentially larger than mean kinetic energies, we may omit the terms containing the spatial differentiation of $\rho_i(\mathbf{r})$. In this approximation,

$$\left(\hat{H}_i[\varphi] + \frac{m_i s_i^2}{2}\right) u_i = \left(E_i + \frac{\hbar^2 k^2}{2m_i}\right) u_i. \quad (6)$$

Then Eqs. (5) reduce to

$$\begin{aligned} \hbar \left(\omega - \mathbf{k} \cdot \mathbf{s}_i - \frac{\hbar k^2}{2m_i}\right) u_i - \sum_j A_{ij} \sqrt{\rho_i \rho_j} (u_j + v_j) &= 0, \\ \hbar \left(\omega - \mathbf{k} \cdot \mathbf{s}_i + \frac{\hbar k^2}{2m_i}\right) v_i + \sum_j A_{ij} \sqrt{\rho_i \rho_j} (u_j + v_j) &= 0. \end{aligned} \quad (7)$$

The latter equations possess a nontrivial solution if the related determinant is zero. To write down the resulting equation in a compact form, we introduce the notation

$$\omega_i^2 \equiv c_i^2 k^2 + \left(\frac{\hbar k^2}{2m_i}\right)^2, \quad c_i^2 \equiv \frac{\rho_i}{m_i} A_i, \quad (8)$$

where $A_i \equiv A_{ii}$. If there would be no interactions between different species, then each component would have the Bogolubov spectrum $\omega_i(k)$ given by Eq. (8). Note that the existence of the Bogolubov spectrum for a single-component Bose-Einstein condensate of trapped atoms is well confirmed by experiment [14]. Define also

$$\omega_{ij} \equiv c_{ij}k, \quad c_{ij}^2 \equiv \left(\frac{\rho_i \rho_j}{m_i m_j} \right)^{1/2} A_{ij}. \quad (9)$$

And let us consider the case of a binary mixture. The spectrum of collective excitations, following from Eq. (7), is given by the equation

$$\left[(\omega - \varepsilon_1)^2 - \omega_1^2 \right] \left[(\omega - \varepsilon_2)^2 - \omega_2^2 \right] = \omega_{12}^4, \quad (10)$$

in which $\varepsilon_i \equiv \mathbf{k} \cdot \mathbf{s}_i$.

It is convenient to connect the system of coordinates with one of the components, say with the first one, so that $\mathbf{s}_1 = 0$. Then $\mathbf{s}_2 \equiv \mathbf{s}$ is the velocity of the second component with respect to the first component. In that case, the spectral equation (10) yields

$$\omega^4 - 2\varepsilon\omega^3 - (\omega_1^2 + \omega_2^2 - \varepsilon^2)\omega^2 + 2\varepsilon\omega_1^2\omega + \omega_1^2(\omega_2^2 - \varepsilon^2) - \omega_{12}^4 = 0, \quad (11)$$

where

$$\varepsilon \equiv \mathbf{k} \cdot \mathbf{s} = ks \cos \vartheta. \quad (12)$$

Equation (11) has four solutions for $\omega = \omega(\mathbf{k})$. But only two of them can be stable, being positively defined for all $\vartheta \in [0, 2\pi]$. The necessary condition for Eq. (11) to possess two positive solutions for arbitrary $\varepsilon \in [-ks, ks]$, by Descartes theorem, is

$$\omega_1^2\omega_2^2 - \omega_{12}^4 > \omega_1^2k^2s^2. \quad (13)$$

In the long-wave limit, when $k \rightarrow 0$, we have the *stability condition*

$$c_1^2c_2^2 - c_{12}^4 > c_1^2s^2. \quad (14)$$

This shows that, when the immovable mixture is stable, satisfying the inequality $c_1^2c_2^2 > c_{12}^4$, it will retain its stability only up to the *critical velocity*

$$s_c = \frac{1}{c_1} \sqrt{c_1^2c_2^2 - c_{12}^4}. \quad (15)$$

Reaching this relative velocity of one component through another, the binary mixture becomes unstable and stratifies onto two spatially separated components. The stratification starts inside the cone of the angle

$$\vartheta_c = \arccos \frac{s_c}{s}. \quad (16)$$

The stability condition (14), with Eqs. (8) and (9), transforms to

$$A_1A_2 - A_{12}^2 > \frac{m_2}{\rho_2} A_1s^2. \quad (17)$$

Respectively, the critical velocity (15) can be written as

$$s_c = \sqrt{\frac{\rho_2(A_1A_2 - A_{12}^2)}{m_2A_1}}. \quad (18)$$

Taking into account the relation between A_{ij} and the scattering length a_{ij} , the stability condition (17) can be presented in the form

$$\frac{a_1a_2}{a_{12}^2} > \frac{(m_1 + m_2)^2}{4m_1m_2} + \frac{m_2^2a_1s^2}{4\pi\hbar^2\rho_2a_{12}^2}. \quad (19)$$

Finally, for colliding wave packets with nonuniform densities $\rho_i(\mathbf{r})$, the stratification would occur not everywhere in space, where the moving packets overlap, but only there, where the inequality

$$\rho_2(\mathbf{r}) > \frac{m_2A_1s^2}{A_1A_2 - A_{12}^2} \quad (20)$$

becomes invalid, provided that $A_1A_2 > A_{12}^2$.

The stability conditions for the mixture of components with relative motion have been obtained here by considering the stability of the spectrum of collective excitations. This is what is called the dynamic stability [2]. There exists as well the notion of thermodynamic stability, when one compares the corresponding free energies, or, at zero temperature, the related internal energies. For immovable mixtures, these two types of stability result in equivalent stability conditions. However, for the mixtures of moving components, these two kinds of stability are not equivalent. To show the latter, we may, for simplicity, employ the uniform approximation, writing the internal energy of the mixture as

$$E_{mix} = \frac{1}{2V} (N_1^2A_1 + N_2^2A_2 + 2N_1N_2A_{12}) + K, \quad (21)$$

where $V = V_1 + V_2$ and the last term is the kinetic energy

$$K = \frac{1}{2} (N_1m_1s_1^2 + N_2m_2s_2^2). \quad (22)$$

If the components are spatially separated, that is, stratified, their total internal energy is

$$E_{str} = \frac{N_1^2A_1}{2V_1} + \frac{N_2^2A_2}{2V_2} + K. \quad (23)$$

Because of the additivity of kinetic energy, the comparison of E_{mix} and E_{str} does not involve the velocities of the moving components, so that the thermodynamic stability condition $E_{mix} < E_{str}$ does not depend on the relative motion of these components. Moreover, employing the thermodynamic stability condition, one commonly invokes the condition of mechanical equilibrium, according to which the pressures of the stratified components,

$$p_i \equiv - \frac{\partial E_{str}}{\partial V_i} = \frac{N_i^2A_i}{2V_i},$$

are equal, so that $p_1 = p_2$. The latter yields the equality

$$\frac{N_1^2}{V_1^2} A_1 = \frac{N_2^2}{V_2^2} A_2 ,$$

from where, recalling that $V = V_1 + V_2$, we have the relations

$$\frac{V}{V_1} = 1 + \sqrt{\frac{N_2^2 A_2}{N_1^2 A_1}} , \quad \frac{V}{V_2} = 1 + \sqrt{\frac{N_1^2 A_1}{N_2^2 A_2}} .$$

This makes it possible to present Eq. (23) as

$$E_{str} = \frac{1}{2V} \left(N_1^2 A_1 + N_2^2 A_2 + 2N_1 N_2 \sqrt{A_1 A_2} \right) + K . \quad (24)$$

Now, comparing Eqs. (21) and (24), one notices that $E_{mix} < E_{str}$ if and only if $\sqrt{A_1 A_2} > A_{12}$. However, the usage of the condition of mechanical equilibrium for moving separated components does not look appropriate. The thermodynamic stability condition can be applied only to immovable mixtures.

The method of studying the stability of mixtures with a relative motion of their components by means of the dynamic stability condition, requiring the positive definiteness for the spectrum of collective excitations, is quite general and can be applied to arbitrary mixtures. In particular, this can also be an atomic-molecular mixture. Such mixtures are formed by means of the Feshbach resonance technique [2,15]. Two Bose atoms can form a quasi-molecule. Then, it is possible to create a two-component mixture consisting of atoms themselves and of the molecules formed by these atoms. When both atoms and molecules are in the Bose-condensed state, they again can be described by the system of the Gross-Pitaevskii equations for the corresponding coherent fields. Thus, if the interaction amplitude for the reaction "atom+atom $\leftrightarrow$ molecule" is denoted by B , then the related Gross-Pitaevskii equations are

$$i\hbar \frac{\partial}{\partial t} \eta_1 = \hat{H}_1[\eta] \eta_1 + B \eta_1^* \eta_2 , \quad i\hbar \frac{\partial}{\partial t} \eta_2 = \hat{H}_2[\eta] \eta_2 + \frac{1}{2} B \eta_1^2 , \quad (25)$$

where $\hat{H}_i[\eta]$ are the same Hamiltonians (2). When atoms as well as molecules are immovable, their mixture is stable under the condition

$$\sqrt{A_1 A_2} > A_{12} + \frac{B}{\sqrt{\rho_2}} ,$$

where $\rho_2 \equiv N_2/V$. But if one is able to organize the relative motion of the atomic and molecular components with respect to each other, then, to find the stability condition for such a moving mixture, one has to follow the same procedure as described above. That is, the form (3) of the coherent fields has to be substituted into Eqs. (25), after which the spectrum of collective excitations can be obtained. Requiring the positiveness of this spectrum yields the stability condition under the relative motion of components.

Concluding, if an immovable mixture of Bose-condensed species is stable, it can stratify if the components move with respect to each other. The stratification occurs when the relative velocity of motion reaches a critical value. The conditions of stability and, respectively, stratification can be derived by linearizing the Gross-Pitaevskii equations for the components, with the inclusion of the Galilean transformation for the coherent fields, as is done in Eq. (3).

References

- [1] Parkins, A.S. and Walls, D.F., 1998, *Phys. Rep.*, **303**, 1.
- [2] Courteille, P.W., Bagnato, V.S., and Yukalov, V.I., 2001, *Laser Phys.*, **11**, 659.
- [3] Fetter, A.L., 2002, *J. Low Temp. Phys.*, **129**, 263.
- [4] Hall, D.S., 2003, *Am. J. Phys.*, **71**, 649.
- [5] Pitaevskii, L.P. and Stringari, S., 2003, *Bose-Einstein Condensation* (Oxford: Oxford University).
- [6] Nepomnyashchy, Y.A., 1974, *Theor. Math. Phys.*, **20**, 904.
- [7] Yukalov, V.I., 1980, *Acta Phys. Pol. A*, **57**, 295.
- [8] Hall, D.S., *et al.*, 1998, *Phys. Rev. Lett.*, **81**, 1539.
- [9] Kozuma, M. *et al.*, 1999, *Phys. Rev. Lett.*, **82**, 871.
- [10] Hagley, E.E. *et al.*, 1999, *Science*, **283**, 1706.
- [11] Maddaloni, P. *et al.*, 2000, *Phys. Rev. Lett.*, **85**, 2413.
- [12] Band, Y.B., Burke, J.P., Simoni, A., and Julienne, P.S., 2001, *Phys. Rev. A*, **64**, 023607.
- [13] Yukalov, V.I., 1998, *Statistical Green's Functions* (Kingston: Queen's University).
- [14] Steinhauer, J., Ozeri, R., Katz, N., and Davidson, N., 2002, *Phys. Rev. Lett.*, **88**, 120407.
- [15] Timmermans, E., Tommasini, P., Hussein, M., and Kerman, A., 1999, *Phys. Rep.*, **315**, 199.

Corresponding author

Prof. V.I. Yukalov

Bogolubov Laboratory of Theoretical Physics

Joint Institute for Nuclear Research

Dubna 141980, Russia

E-mail: yukalov@thsun1.jinr.ru

Tel: 7 (096) 216-3947 (office); 7 (096) 213-3824 (home)

Fax: 7 (096) 216-5084